

The Ability of Variant Peptides to Reverse the Nonresponsiveness of T Lymphocytes to the Wild-Type Sequence p53_{264–272} Epitope¹

Thomas K. Hoffmann,* Douglas J. Loftus,[§] Koji Nakano,* Markus J. Maeurer,[¶] Kazuaki Chikamatsu,* Ettore Appella,[§] Theresa L. Whiteside,*^{†‡} and Albert B. DeLeo^{2*†}

Recently, we observed that CTL specific for the wild-type (wt) sequence p53_{264–272} peptide could only be expanded ex vivo from PBMC of a subset of the HLA-A2.1⁺ normal donors or cancer patients tested. Surprisingly, the tumors of the responsive patients expressed normal levels of wt p53 and could be considered unlikely to present this epitope. In contrast, tumors of nonresponsive patients accumulated mutant p53 and were more likely to present this epitope. We sought to increase the responsive rate to the wt p53_{264–272} peptide of PBMC obtained from normal donors and patients by identifying more immunogenic variants of this peptide. Two such variants were generated by amino acid exchanges at positions 6 (6T) and 7 (7W) of the peptide. These variants were capable of inducing T cells from PBMC of nonresponsive donors that recognized the parental peptide either pulsed onto target cells or naturally presented by tumors. TCR V β analysis of two T cell lines isolated from bulk populations of effectors reactive against the wt p53_{264–272} peptide, using either the parental or the 7W variant peptide, indicated that these T cells were expressing identical TCR V β 13.6/complementarity-determining region 3/J region sequences. This finding confirms the heteroclitic nature of at least one of the variant peptides identified in this study. The use of variant peptides of the wt p53_{264–272} epitope represents a promising approach to overcoming the nonresponsiveness of certain cancer patients to this self epitope, thereby enhancing its potential use in tumor vaccines for appropriately selected cancer patients. *The Journal of Immunology*, 2002, 168: 1338–1347.

Genetic alterations in p53 occur in a wide range of human tumors, including oral squamous cell carcinomas (OSCC)³ (1). The most common type of genetic alteration in p53 involves a missense mutation that is usually accompanied by accumulation of the altered molecules in the cytosol of tumor cells. Initially, the effort to develop p53-based vaccines focused on these missense mutations, which are tumor specific in nature. However, missense mutations have limited clinical usefulness, because of the requirement that they occur within or create epitopes that could be presented by MHC molecules expressed by the individual patient. On the other hand, the majority of p53 epitopes derived from these altered p53 molecules would be wild type in sequence, representing a new class of tumor-associated self

Aggs that are candidates for use in the development of broadly applicable cancer vaccines (1–5).

To date, five MHC class I-restricted, naturally presented human wild-type (wt) sequence p53 epitopes have been identified. They have been shown to be able to induce epitope-specific CTL from PBMC obtained from healthy individuals (1, 6–11). The p53_{125–134} epitope is HLA-A24 restricted (11), while the other four, p53_{65–73}, p53_{149–157}, p53_{217–225}, and p53_{264–272}, are HLA-A2.1 restricted. Among these, the wt p53_{264–272} peptide has been the most intensively investigated (1, 2, 6–8, 12).

The potential of wt p53 epitopes as targets for immunotherapy, however, remains uncertain due to the several critical concerns related to immunological recognition of this truly self tumor Ag. Using HLA-2.1-transgenic wt (p53^{+/+}) and p53^{null} (p53^{-/-}) mice, Sherman and colleagues (13–15) have demonstrated that the CTL repertoire available for wt p53 self epitopes in p53^{+/+} mice is limited to intermediate affinity T cells, because the higher affinity CTL are either deleted or tolerized. Apparently, this situation occurs in humans as well, as only CTL with intermediate affinity for the wt p53_{264–272} epitope have been generated to date from PBMC obtained from normal donors as well as cancer patients (7, 12). This observation raises the question of whether such CTL are potent enough to be effective in tumor eradication.

Another concern relates to our experience that PBMC obtained from only some HLA-A2.1⁺ healthy donors and patients with OSCC were responsive to in vitro stimulation (IVS) with the wt p53_{264–272} peptide pulsed onto autologous dendritic cells (DC) (7, 12). Furthermore, CTL reactive against this epitope could only be generated from T cell precursors in PBMC of patients whose tumors were not likely to present this epitope. The analysis of these

*University of Pittsburgh Cancer Institute and Departments of [†]Pathology and [‡]Otolaryngology, University of Pittsburgh School of Medicine, Pittsburgh, PA 15213; [§]National Cancer Institute, Bethesda, MD 20892; and [¶]Department of Medical Microbiology, Johannes Gutenberg University, Mainz, Germany

Received for publication August 7, 2001. Accepted for publication November 27, 2001.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported in part by National Institutes of Health Grant PO1-DE12321 (to T.L.W. and A.B.D.) and Department of Defense Grant BC995591 (A.B.D.). T.K.H. was supported by a postdoctoral fellowship from the Dr. Mildred Scheel Stiftung für Krebsforschung.

² Address correspondence and reprint requests to Dr. Albert B. DeLeo, University of Pittsburgh Cancer Institute, University of Pittsburgh School of Medicine, Division of Basic Research, Biomedical Science Tower W956, 211 Lothrop Street, Pittsburgh, PA 15213. E-mail address: deleo@imap.pitt.edu

³ Abbreviations used in this paper: OSCC, oral squamous cell carcinoma; APL, altered peptide ligand; CDR, complementarity-determining region; DC, dendritic cell; IVS, in vitro stimulation; wt, wild type.

tumors indicated no accumulation of p53 or accumulation of mutant p53 with a missense mutation at codon 273, a site known to block processing of the wt p53_{264–272} epitope (16). In contrast, PBMC obtained from patients with tumors considered capable of presenting the wt p53_{264–272} epitope (i.e., tumors that accumulate mutant p53) were nonresponsive to IVS with wt p53_{264–272}-pulsed autologous DC. These findings have led us to conclude that CTL specific for the wt p53_{264–272} epitope might play a role in the outgrowth of epitope-loss tumor cells, which are able to escape from the host immune system. This conclusion was further strengthened by the results of a recently completed study in our laboratories that used tetrameric peptide/MHC class I complexes to determine frequencies and characteristics of the p53_{264–272}-specific CTL in unstimulated PBMC obtained from 30 OSCC patients and 31 normal donors (data not shown).⁴

Because these observations suggest that it may be possible to accurately predict *ex vivo* the responsiveness of cancer patients to immunotherapy targeting this epitope, we felt that a means of circumventing the nonresponsiveness of individuals needed to be investigated to proceed with the development of wt p53-based vaccines. One solution is to identify a heteroclitic peptide or, in more precise terms, an altered peptide ligand (APL) with enhanced functional activity relative to the parental wt p53_{264–272} peptide. By substituting amino acids at various positions of an epitope that contact MHC class I and/or TCR, an array of APL with biological potencies higher than those of the parental epitopes has been identified for various antigenic determinants (17–26). In applying this strategy to the wt p53_{264–272} epitope, we anticipated that an APL might induce CTL-mediated responses that cross-react with the parental epitope and that these CTLs also might demonstrate enhanced avidities relative to CTLs induced by the parental peptide. Most importantly, we sought to determine whether an APL would be able to induce anti-wt p53_{264–272} CTL from PBMC that were nonresponsive to the parental peptide, particularly the PBMC obtained from patients whose tumors accumulate mutant p53 and are considered to have the potential to present this epitope.

Materials and Methods

Cell lines and cell culture

The following HLA-A2⁺ OSCC cell lines were obtained from the American Type Culture Collection (ATCC, Manassas, VA): SCC-4 and SCC-9. The SCC-4 cell line expresses and accumulates p53 expressing a missense mutation at codon 151 but does not present the wt p53_{264–272} epitope (6). The SCC-9 cell line expresses an altered p53 molecule with a deletion of codons 274–285. It does not accumulate p53 molecules, yet presents the wt p53_{264–272} epitope. In addition, the p53^{mut} osteosarcoma cell line, SaOS-2, was obtained from ATCC. The cloned p53⁺ cell line, SaOS-2Cl3, was derived by transduction of SaOS-2 cells with a p53 cDNA expressing a missense mutation in codon 143 (7). The HLA-A2⁺ OSCC cell line PCI-13 has been described previously (27). It expresses a p53 missense mutation in codon 286 (Glu to Lys) and presents the wt p53_{264–272} epitope. Tumor cells were cultured in plastic culture flasks (Costar, Cambridge, CA) under standard conditions (37°C, 5% CO₂ in a fully humidified atmosphere) in complete medium consisting of DMEM (Life Technologies, Grand Island, NY) supplemented with 10% (v/v) heat-inactivated FBS, 2 mM L-glutamine, 50 µg/ml streptomycin, and 50 IU/ml penicillin (all from Life Technologies). The T2 cell line was also obtained from ATCC and maintained in RPMI 1640 (Life Technologies) containing 10% heat-inactivated FBS, 2 mM L-glutamine, and antibiotics. The cultures were routinely tested and found to be free of mycoplasma contamination (Gen-Probe, San Diego, CA).

Peptides

The CTL-defined, HLA-A2.1-binding peptide, LLGRNSFEV (1), corresponding to wt p53_{264–272}, as well as single amino acid exchange variants of this peptide were synthesized by standard *N*-(9-fluorenyl)methoxycarbonyl methology. Peptides were purified by reversed-phase HPLC, and their amino acid sequence was confirmed by mass spectrometry analysis. All peptides were dissolved in DMSO (Fisher Scientific, Pittsburgh, PA) at 1 mg/ml and diluted with PBS just before use. The 19 variant peptides contain single amino acid exchanges with a bias toward retention of a high degree of similarity to the central region of the parental peptide. The variant peptides are designated 1E, 1F, 1V, 3L, 3F, 3W, 4K, 4L, 5K, 5L, 6G, 6T, 6Y, 7L, 7P, 7Y, 7W, 8A, and 8Y, in which numbers denote the position within the parental sequence and letters refer to exchanged amino acids.

MHC stabilization assay

T2 cells were incubated overnight at room temperature before use in this assay. Cells were washed and incubated at a cell density of $2 \times 10^5/0.2$ ml of complete medium with various peptides at final concentrations of 1×10^{-5} – 1×10^{-10} M for 3 h at room temperature, followed by a 3-h incubation period at 37°C. After washing with PBS, cells were incubated at 4°C for 30 min with anti-HLA class I mAb, W6/32 (HB95; ATCC), and then with FITC-conjugated goat anti-mouse Ig (Caltag Laboratories, Burlingame, CA) as a secondary Ab. Fluorescence of viable T2 cells was measured at 488 nm in a FACScan flow cytometer (BD Biosciences, San Jose, CA), and the level of MHC class I expression was determined by evaluating the mean fluorescence intensity of stained T2 cells. Cells incubated either at room temperature or 37°C in the absence of peptide served as controls.

Generation of anti-p53 CTL with peptide-pulsed autologous DC

Peripheral blood or leukapheresis products were obtained from previously studied HLA-A2.1⁺ individuals: seven normal donors and six OSCC patients (12). PBMC were isolated by sedimentation over Ficoll-Hypaque (Amersham Pharmacia Biotech, Piscataway, NJ). The study was approved by the Institutional Review Board at the University of Pittsburgh, and written informed consent was obtained from each individual donating peripheral blood. PBMC were phenotyped for HLA-A2 expression by flow cytometry, using anti-HLA-A2 mAb, BB7.2 (HB82; ATCC), and a mouse IgG isotype as a control. The verification of the A0201 subtype was performed using PCR with sequence-specific primers, as previously described (12).

Human DC were generated from PBMC according to a modification of the method of Sallusto and Lanzavecchia (28), as described by us earlier (7). DC were harvested on day 6, phenotyped by flow cytometry, and then resuspended in AIM-V medium (2×10^6 cells/ml) containing 10 µg/ml peptide and incubated at 37°C for 4 h. The peptide-pulsed DC were then cocultured with autologous PBMC in 24-well tissue culture plates (Costar) in a final volume of 2 ml/well AIM-V medium supplemented with 10% (v/v) human AB serum (Pel-Freez Biologicals, Brown Deer, WI) and 25 ng/ml IL-7 (Genzyme, Cambridge, MA) for the first 72 h and, additionally, with 20 IU/ml IL-2 (Chiron-Cetus, Emeryville, CA) for the remaining time in culture. The lymphocytes were restimulated 1 wk later with peptide-pulsed autologous DC. Irradiated (3000 rad) autologous PBMC were used as APC after the third round of restimulations. Microcultures of CTL lines recognizing the wt p53_{264–272} or 7W peptide were isolated from bulk populations of effectors by limiting dilution (1 cell/well/96-well plates), and the lines were maintained in cytokine-supplemented media plus peptide-pulsed APC, as previously described (7). Specificities of generated T cells were determined using one or more of a panel of assays detailed below. The TCR Vβ expression on T cells in bulk CTL populations and cell lines derived from them was done using the IOTest β Mark TCR Vβ Repertoire kit (Beckman Coulter, San Diego, CA).

ELISPOT assay for IFN-γ

The ELISPOT assay was performed in 96-well plates with nitrocellulose membrane inserts (Millipore, Bedford, MA), as previously described by us (29). The capture and detection anti-IFN-γ mAb were purchased from Mabtech (Nacka, Sweden). The spots were counted by computer-assisted image analysis (ELISPOT 4.14.3; Zeiss, Jena, Germany). For Ab-blocking experiments, target cells were preincubated with anti-HLA class I mAb for 30 min. Cryopreserved aliquots of PBMC obtained from a normal donor were thawed and, after stimulation with PMA (1 ng/ml) and ionomycin (1 µM; both from Sigma-Aldrich, St. Louis, MO), were used as a positive control for each assay.

⁴ T. K. Hoffmann, A. Donnenberg, S. Finkelstein, K. Chikamatsu, V. Donnenberg, U. Friebe, E. Appella, A. B. DeLeo, and T. L. Whiteside. Frequencies of tetramer⁺ T cells specific for the wild type sequence p53_{264–272} peptide in the circulation of patients with head and neck cancer. *Submitted for publication.*

The interassay reproducibility of the assay was acceptable with a coefficient of variation = 15% ($n = 30$).

Cytotoxicity assay

The 4-h ⁵¹Cr release assay was performed at various E:T ratios, as previously described (7). Briefly, sensitized targets were labeled with ⁵¹Cr for 45 min at 37°C, washed, and added to wells of 96-well plates (1×10^5 cells/well). Effector T cells were then added to give various E:T ratios. When Ab-blocking experiments were performed, target cells were incubated with anti-HLA class I mAb or the anti-HLA-DR mAb, L243 (HB55; ATCC), for 30 min before adding effector cells. The percentage of specific lysis was calculated according to the formula:

$$\% \text{ specific lysis} = \frac{\text{experimental cpm} - \text{control cpm}}{\text{maximal cpm} - \text{control cpm}} \times 100$$

Flow cytometry analysis using HLA-A2.1/peptide tetrameric complexes (tetramer)

The streptavidin-PE-labeled tetramers used in this study were obtained from the tetramer core facility of the National Institute of Allergy and Infectious Disease (Atlanta, GA). Three-color flow cytometry assays (FACScan; BD Biosciences) were performed with PerCP anti-CD3, FITC anti-CD8, and PE-tetramer. The specificity of the HLA-A2.1/p53₂₆₄₋₂₇₂ tetramer was confirmed by its staining of a CTL line specific for this p53 epitope and by the lack of staining of irrelevant CTL or HLA-A2⁻ PBMC of healthy donors, as previously described (30). The additional PE-conjugated HLA-A2.1/tetramer used in this study contained the 7W variant peptide. Generally, 75,000 events per sample were collected progressively after live gating on lymphocytes by forward and side scatter.

TCR and CDR3 spectratyping

RNA was extracted from p53₂₆₄₋₂₇₂-specific CTL lines generated using parental or the 7W variant peptide, followed by reverse transcription into cDNA, as previously described (31). Screening for expression of TCR V chains was performed using the primers described by Puisieux et al. (31) for TCR V β amplification, followed by a runoff reaction with fluorophore-labeled primers specific for the C region of the TCR β (5'-TGTGCAC CTCCTTCCCATTCACC) chain. Labeled runoff products were subjected to DNA fragment analysis, as described (32). Finally, amplified products were directly subjected to DNA sequence analysis using ABI 310 sequencer (PerkinElmer, Weiterstadt, Germany).

Statistical analysis

A two-tailed Wilcoxon rank sum test was performed to analyze ELISPOT data. An unpaired two-tailed Student's *t* test was used to interpret differences in CTL reactivities against different target cells and in the presence of blocking Ab in cytotoxic assays, and differences between the number of spots obtained from T cells incubated with T2 cells pulsed with relevant p53 peptides vs that obtained using T2 cells pulsed with the irrelevant

gp100 peptide in ELISPOT assays. Differences were considered significant when $p < 0.05$.

Results

Selection of variant p53₂₆₄₋₂₇₂ peptides recognized by anti-p53₂₆₄₋₂₇₂ CTL

Because the parental peptide binds efficiently to HLA-A2.1 molecules, all of the APL considered in this study represent single amino acid exchanges at nonanchor residues for the purpose of enhancing the interactions of the variant peptides with the TCR rather than MHC class I molecules. Nineteen variants of the wt p53₂₆₄₋₂₇₂ peptide were screened for their recognition by a bulk population of anti-wt p53₂₆₄₋₂₇₂-specific CTL that was maintained in our laboratory (7). T2 cells pulsed with the individual peptides at a fixed concentration of 1×10^{-6} M peptide served as targets for these CTL in a ⁵¹Cr release cytotoxic assay. Significant cytotoxic reactivity against T2 cells pulsed with three of the 19 variant peptides, namely, 6T, 7W, and 7P, was detected (data not shown). Therefore, these three variant peptides were selected for further characterization.

Variant peptide binding to HLA-A2.1 molecules

Binding of the 6T, 7W, and 7P variant peptides to HLA-A2.1 molecules was compared with that of the parental peptide in an MHC stabilization assay. The relative mean fluorescence intensity of parental and variant peptide-stabilized HLA-A2 molecules on T2 cells is shown in Fig. 1. All the peptides showed stabilization of HLA-A2 molecules in a dose-dependent manner within the concentration range of 1×10^{-5} – 1×10^{-9} M. However, in general, the binding affinities of the variant peptides to HLA-A2.1 molecules on T2 cells were slightly lower than that of the parental wt peptide ($\text{wt} > 6\text{T} \geq 7\text{W} > 7\text{P}$).

Affinity of p53₂₆₄₋₂₇₂-specific CTL for variant peptides

The affinity of the bulk population of anti-p53₂₆₄₋₂₇₂-specific CTL for the variant peptides was determined in a 4-h ⁵¹Cr release assay using T2 cells pulsed with these peptides at concentrations ranging from 1×10^{-5} to 1×10^{-12} M as target cells. As shown in Fig. 2, at concentrations $< 1 \times 10^{-8}$ M, the dose-response curves of the three variant peptides were shifted to the left relative to that of the parental wt peptide. Because the increased responsiveness of the CTL for these variant peptides cannot be attributed to enhanced binding to HLA-A2.1 molecules, these results are consistent with an increased affinity of TCR for the variant peptides.

FIGURE 1. Identification of three HLA-A2.1-binding variant peptides of the wt p53₂₆₄₋₂₇₂ epitope. T2 cells were incubated with parental p53₂₆₄₋₂₇₂ peptide (LLGRNSFEV) or 6T, 7P, or 7W variant peptides at final concentrations of 1×10^{-5} – 1×10^{-10} M. The relative mean fluorescence intensities of FITC-conjugated anti-MHC class I mAb (W6/32) are indicative of peptide-stabilized MHC class I molecules on T2 cells.





FIGURE 2. Variant peptides are recognized by anti-wt p53_{264–272}-specific CTL line. T2 cells were pulsed with different peptide concentrations and tested as targets in a 4-h ⁵¹Cr release assay at the E:T ratio of 10:1.



Characterization of wt p53_{264–272}-specific CTL generated from PBMC obtained from normal donors using variant peptides

Previously, we reported that CTL reactive against the wt p53_{264–272} epitope could be generated from PBMC obtained from only two of the seven HLA-A2.1⁺ normal donors tested (12). Analyses involving multiple cryopreserved samples derived from leukopaks obtained from two of the normal donors (a responder and a nonresponder) confirmed the consistency of responses of these donors' PBMC to the parental peptide. In the same experiments in which the seven donors' PBMC were tested for induction using the parental peptide, CTL reactive against this peptide could be generated from five of these seven PBMC using either the 6T or 7W variant peptide. Included in this group were three nonresponsive PBMC (Table I); PBMC obtained from donors 6 and 7 responded to the 7W variant, while PBMC obtained from donor 4 responded to the 6T peptide. None of the seven PBMC tested responded to the 7P variant peptide.

The bulk populations of variant-induced cells generated from PBMC obtained from donors 6 and 7 effectively recognized and lysed T2 cells pulsed with the parental peptide in ELISPOT for IFN- γ and cytotoxicity assays. Fig. 3, A and B, shows the results obtained with the effectors generated from PBMC obtained from donor 7. Unpulsed T2 target cells or T2 cells pulsed with an

relevant HLA-A2.1-binding peptide, the melanoma-associated gp100 peptide (33), were not recognized by these CTL in either assay to any noticeable extent. More importantly, these effector cells were also capable of recognizing the naturally presented epitope, as evidenced by their ability to lyse PCI-13 and SCC-9 tumor cells as well as mutant p53-transfected SaOS-2 Cl3 cell lines in a MHC class I-restricted manner (Fig. 3C). No significant cytotoxicity was noted against HLA-A2⁺ tumor cell lines, SCC-4 and SaOS-2, which do not present the epitope.

The CTL generated from PBMC obtained from donor 4 using the 6T peptide yielded effectors with reactivity comparable with that of the 7W variant-induced CTL. The 6T-induced T cells were responsive to wt p53_{264–272}-pulsed T2 cells in the ELISPOT for IFN- γ assay (Fig. 4A), and cytolytic against the OSCC lines, SCC-9 and PCI-13, as well as SaOS-2Cl3 (Fig. 4B). This response was blocked by anti-HLA class I mAb but not anti-HLA-DR mAb. No significant reactivity was obtained against the tumor cell line SCC-4. The reactivity of these effectors against SaOS-2 cells in the analysis shown in Fig. 4B was higher than normally detected against this p53^{null} cell line, using bulk populations of anti-p53 effectors (7, 12). However, the reactivity of the 6T-induced effectors against SaOS-2 targets was not significantly blocked by anti-HLA class I mAb and thus could be attributed to nonspecific effectors present in the bulk population. In summary, variant-induced effector T cells had similar reactivities against the parental epitope as those reported previously for the parental peptide-induced effectors from responsive normal donors as well as OSCC patients (7, 12).

Table I. Summary of the anti-p53 CTL responses of PBMC obtained from normal donors following IVS using variant p53_{264–272} peptides^a

Donor	Anti-p53 CTL Response After IVS With			
	wt p53 _{264–272}	Variant 6T	Variant 7P	Variant 7W
1	+	–	–	+
2	+	–	–	+
3	–	–	–	–
4	–	+	–	–
5	–	–	–	–
6	–	–	–	+
7	–	–	–	+

^a PBMC were stimulated with the peptide-pulsed autologous DC or PBMC in two to four IVS cycles. Effector cell reactivity was tested in ELISPOT and cytotoxicity assays; – indicates that no specific reactivity against the wt p53_{264–272} peptide was observed, while + indicates that effectors were reactive against variant and parental peptides. Results using the parental wt p53_{264–272} were reported in a previous publication from this laboratory (12).

Characterization of wt p53_{264–272}-specific CTL generated from PBMC of a nonresponsive OSCC patient using a variant peptide

The critical test of the variant peptides was whether their use could induce CTL capable of recognizing the anti-wt p53_{264–272} epitope from nonresponsive patients whose tumors were considered capable of presenting this epitope (12). The nonresponsiveness of PBMC obtained from at least one of these donors, patient 3, has been repeatedly confirmed during the past 2 years using blood samples obtained at different times, as well as multiple cryopreserved leukapheresis samples obtained from this patient. As shown in Table II, none of the PBMC from three of these patients responded to the 6T or 7P variant peptides. However, the 7W variant




FIGURE 3. The 7W variant peptide-induced effectors induced from PBMC obtained from a nonresponsive healthy donor recognize the parental wt p53₂₆₄₋₂₇₂ peptide pulsed onto target cells or naturally presented by tumors. *A*, Recognition of peptide-pulsed T2 cells in ELISPOT for IFN- γ assays. Effectors were tested against T2 cells pulsed with an irrelevant gp100 peptide, the wt p53₂₆₄₋₂₇₂ peptide, or the variant peptides at 10 μ g/ml. *B*, Lysis of T2 cells pulsed with various peptides at an E:T of 1:10. *C*, Lysis of tumor targets naturally presenting the epitope at an E:T ratio of 40:1 in the presence or absence of anti-MHC class I or anti-HLA-DR mAb. *, A significant ($p < 0.05$) difference relative to IgG controls.

peptide did induce the ex vivo generation of anti-wt p53₂₆₄₋₂₇₂ CTL from PBMC of patient 3, whose autologous tumor, PCI-13, presents this epitope (7, 12). The affinity of these effectors for the parental epitope was comparable with that of a bulk population of CTL induced using the parental epitope, and was in the range of 1×10^{-9} M (Figs. 2 and 5A). Furthermore, the 7W-induced CTL were cytotoxic against a panel of tumor cell lines naturally presenting the wt p53₂₆₄₋₂₇₂ epitope, including the autologous PCI-13 cell line, and this reactivity was MHC class I restricted (Fig. 5B). This result clearly illustrates the potential value of the 7W variant




FIGURE 4. The 6T variant peptide-induced effectors induced from PBMC obtained from nonresponsive healthy donor recognize the parental wt p53₂₆₄₋₂₇₂ epitope pulsed onto target cells or naturally presented by tumors. *A*, Recognition of peptide-pulsed T2 cells in ELISPOT assays. Effectors were tested against T2 cells pulsed with either an irrelevant gp100 peptide, the wt p53₂₆₄₋₂₇₂ peptide, or a variant peptide at 10 μ g/ml. *B*, Lysis of tumor targets naturally presenting the epitope at an E:T of 40:1 in the presence or absence of anti-MHC class I or anti-HLA-DR mAb. *, A significant ($p < 0.05$) difference relative to IgG controls.

peptide in immunotherapy targeting the wt p53₂₆₄₋₂₇₂ epitope in individuals like OSCC patient 3.

Tetramer-binding and TCR V β usage by T cell microcultures reactive against parental and/or variant peptides

The ability of the variant peptides to induce the generation of CTL specific for wt p53₂₆₄₋₂₇₂ from nonresponder PBMC raised the question of the relationship between these CTL and those induced by the parental peptide in responder PBMC. The need to investigate this relationship became evident when the cross-reactive bulk population of CD8⁺ T cells induced with the 7W variant peptide from normal donor 7 was stained with the parental or variant tetramer. Whereas only $\sim 2\%$ tetramer^{dim} cells were detected with the parental tetramer, a cluster of $\sim 40\%$ tetramer^{bright} cells was detected with the 7W tetramer (Fig. 6, A and B). One possible explanation for this observed difference was that the variant peptide induced a single CD8⁺ T cell population that bound the variant tetramer with higher avidity/stability than did the parental tetramer. Another possible explanation was that the variant peptide induced two distinct populations of CD8⁺ T cells; one was cross-reactive and bound both tetramers (most likely with different avidities), while the other was specific for the 7W variant and bound the 7W tetramer with high avidity. The two possibilities could be

Table II. Summary of the anti-p53 CTL responses of PBMC obtained from nonresponsive OSCC patients following IVS using variant p53₂₆₄₋₂₇₂ peptides

Patient	Tumor ^a		Anti-p53 CTL Response After IVS ^b		
	p53 genotype	p53 protein	Variant 6T	Variant 7P	Variant 7W
1	Mutant R248W	+	-	-	-
2	Mutant V 157 F	+	-	-	-
3	Mutant E286K	+	-	-	+

^a Patients' tumors were analyzed for genetic alterations in p53 exons 5-8, and the identified codon and missense mutations are denoted. The level of p53 expression in tumors was determined by immunohistochemistry, using anti-p53 mAb, and + denotes accumulation of p53 (12).

^b PBMC were stimulated with the peptide-pulsed autologous DC or PBMC in two to four IVS cycles. Effector cell reactivity was tested in ELISPOT and cytotoxicity assays; - indicates that no specific reactivity against the wt p53₂₆₄₋₂₇₂ peptide was observed, while + indicates that effectors were reactive against variant and parental peptides.

distinguished based on TCR usage of the T cells involved in recognition of these peptides. To accomplish this, T cell microcultures were established by limiting dilution from bulk CTL populations induced with either parental or variant peptide. Several T cell clones from each type of microculture were expanded for further analysis. Based in part on their rates of proliferation as well as

peptide specificities (Fig. 7), four oligoclonal T cell lines, designated 2, 4, 53, and 68, were selected for TCR analysis by complementarity-determining region (CDR)3 spectratyping.

Two of the cell lines analyzed, 53 and 68, were derived from the bulk population of 7W-induced CTL that was described above and shown to exhibit differential staining with the parental and 7W

A. ELISPOT





FIGURE 5. The 7W variant peptide-induced effectors obtained from a nonresponsive OSCC patient recognize the parental wt p53₂₆₄₋₂₇₂ peptide pulsed onto target cells or naturally presented by tumors, including PCI-13, the autologous tumor cell line. A, Affinity of effectors for the parental wt p53₂₆₄₋₂₇₂ peptide as determined in ELISPOT assay. T2 cells were incubated with parental peptide at concentrations of 1 × 10⁻⁶–1 × 10⁻¹⁰ M. T2 cells pulsed with an irrelevant gp100 peptide served as a control. B, Lysis of tumor targets naturally presenting the epitope at an E:T of 40:1 in the presence or absence of anti-MHC class I or anti-HLA-DR mAb. *, A significant (*p* < 0.05) difference relative to IgG controls.

B. ⁵¹Cr-Release Assay






FIGURE 6. CD8⁺ cells induced from PBMC of a nonresponsive normal donor using the 7W variant peptide that recognizes the parental peptide express TCR V β 13.6. Three-color flow cytometry analysis of CD8⁺ cells stained with HLA-A2.1 tetramers containing either the parental peptide (A), 7W variant peptide (B), or the 7W tetramer and anti-V β 13.6 mAb (C). The numbers in the *upper right quadrants* indicate the percentage of tetramer⁺ cells. The analyses shown in A and B involved 75,000 events, while in C 10,000 events were analyzed.

tetramers (Fig. 6). Although tetramer analysis of the bulk population suggested that the vast majority of CTL were 7W specific, one of the several T cell clones isolated was cross-reactive. The 53 cell line, which was specific for the 7W variant, was found to express V β 9. The 68 cell line, which recognized the variant as well as the parental peptide, was found to express V β 13.6 with completely different CDR3 and J regions from those expressed by line 53 (Table III). Consistent with these findings was the result of a combined tetramer/V β expression flow analysis of the bulk population from which the 53 and 68 cell lines were derived. We observed that the ~40% cells that stained as a distinct cluster with the 7W tetramer were V β 13.6⁻, whereas the ~2% cells that stained weakly with this tetramer were V β 13.6⁺ (Fig. 6C).

The other two cell lines analyzed, 2 and 4, were derived from bulk populations of effectors induced with the parental peptide from PBMC obtained from a patient (patient 2 in Ref. 12) and a normal donor (donor 2, Table I), respectively. The 2 cell line was found to express V β 13.6, with identical motifs for the CDR3 and J regions as the 68 cell line, which was induced from a different individual using the 7W peptide (Table III). The observation that cross-reactive T cells induced by the variant peptide from PBMC obtained from one donor express the identical TCR/CDR3/J region as that expressed by T cells induced with the parental peptide from another donor illustrates the heteroclitic nature of the 7W variant peptide.

The 4 cell line was shown to express V β 1 with a CDR3 and J region sequence distinct from those of any of the other cell lines analyzed (Table III). Interestingly, V β analysis of the bulk popu-

lation of effectors from which the 4 cell line was isolated detected mainly V β 1 and V β 13 CD8⁺ cells. Another bulk population of parental peptide-induced CTL, which was obtained from a different normal donor (7) and used to identify 7W and 6T peptides as potential APLs (Table II), was found to consist of >90% V β 1⁺ cells (data not shown). These results, summarized in Table IV, are strongly suggestive of a relatively limited TCR V β usage being involved in recognition by CTL of the HLA-A2.1-restricted, wt p53_{264–272} epitope, regardless of whether these cells are induced by the parental or variant peptide.

Discussion

Most studies of APL of tumor Ags involve amino acid exchanges at anchor residue positions of the peptide to enhance its binding to class I MHC molecules. The recent modification of the HLA-A2.1-restricted wt p53_{149–157} peptide at anchor position 2 to improve its binding to the restriction element and immunogenicity is one example of this approach (26). Particularly relevant to this study are variants designed to enhance TCR/peptide interactions rather than increase MHC binding, such as the HLA-A2.1-restricted, melanoma-associated MART1/Melan A_{27–35} and carcinoembryonic Ag, CAP1, peptides, which involve amino acid exchanges in residues other than anchor positions (18–21). While the binding affinities of these variant and parental peptides to HLA-A2.1 molecules are comparable, amino acid exchanges of these peptides at nonanchor positions yielded variant peptides that were more immunogenic than the parental peptides.

Since the parental wt p53_{264–272} peptide has a reasonable affinity for HLA-A2.1 molecules ($>1 \times 10^{-9}$ M), the 19 p53_{264–272} variants designed for this study had unmodified anchor positions. Among the amino acid exchanges tested, those at position 6 (6T) and position 7 (7W) appeared to be promising. Since both variants have lower affinities than the parental peptide for HLA-A2.1 molecules, their ability to increase the frequency of anti-p53_{264–272} CTL responses generated from nonresponsive PBMC does not appear to be due to their enhanced binding to HLA-A2.1 molecules. Instead, their increased immunogenicity might be due to the replacement or counterbalancing of residues causing adverse TCR-peptide interactions. Such a replacement could result in an improved interaction of the peptide/MHC complex with TCR and a subsequent expansion of T cells capable of recognizing the parental epitope (23, 24). Two lines of evidence support this conclusion. First, using the parental tetramer to determine the frequency of tetramer⁺ precursor T cells in unstimulated PBMC obtained from normal donors and patients with cancer, we found that most of the nonresponsive individuals had markedly lower frequencies of these cells in their peripheral circulation than did the responders (data not shown).⁴ Second, the parental and variant peptides were found to engage and expand T cells expressing the same TCR in PBMC obtained from responsive and nonresponsive donors (see Tables III and IV). These findings support the concept that increased stability of interaction with the TCR is the basis for the enhanced functional activity of the 7W variant peptide.

Although the use of variant peptides did reverse the nonresponsiveness in IVS of PBMC obtained from some donors, their use did not yield high-affinity CTL. The persistence of low-affinity CTL against self tumor peptides, such as wt p53 epitopes, which is considered a true consequence of tolerance (15), might be due to a limited TCR repertoire being available for recognition of these epitopes. Our analyses detected the predominant use of only two TCR V β families, V β 1 and V β 13.6, being involved in CTL recognition of the wt p53_{264–272} epitope in four different donors. Furthermore, in two different donors, identical usage by the parental




FIGURE 7. Specificity of T cell lines analyzed for TCR V β usage. Lines 2 and 4 were derived from bulk populations of effectors induced from an OSCC patient and a normal control, respectively, using the parental peptide. Both are cross-reactive against the variant peptide. Lines 53 and 68 were derived from a bulk population of effectors induced from a nonresponsive healthy donor with the 7W variant peptide. The 53 cell line is specific for the 7W variant, while the 68 cell line is cross-reactive.

and variant peptide was detected. In contrast, an analysis of responses in HLA-A2.1⁺ patients to repeated immunizations with an anchor position-variant peptide of the melanoma-associated gp100₂₀₉₋₂₁₇ epitope demonstrated that the appearance of higher-affinity T cells was associated with an expansion of the TCR repertoire rather than an increased oligoclonal response (33). In the future, additional data on TCR usage of cross-reactive and variant-specific CTL cell could allow for extensive molecular modeling of the interactions within the trimeric complexes and, perhaps, the design of APL with more enhancing properties than those of the 6T and 7W variants. These variants might engage more diverse populations of T cells that are capable of cross-recognition of the parental epitope with, perhaps, higher avidity. However, the apparent outgrowth of epitope-loss tumors in OSCC patients responsive to this epitope suggests that even intermediate-affinity CTL recognizing wt p53₂₆₄₋₂₇₂ might be effective in tumor eradication (12).

To fully estimate the potential of p53-based vaccines in immunotherapy of cancer, it is becoming increasingly apparent that an array of T cell-defined wt p53 epitopes needs to be analyzed, and strategies for optimal induction of T cells recognizing these epitopes need to be further evaluated. In this regard, the use of genetically modified DC expressing intact wt p53 appears to enhance the generation and increase the frequency of antitumor effectors from PBMC of normal donors and cancer patients (34). The p53-based immunotherapy also might be critically dependent on targeting the right epitopes and matching a patient's ability to respond ex vivo to wt p53 epitopes with the potential of his/her tumor to present these epitopes for immune recognition. Again, of course, it is necessary to be aware that a patient's ex vivo responsiveness to these epitopes does not guarantee a successful in vivo response to immunization with them. In this study, HLA-A2.1⁺ patient 3 with OSCC, for whom the tumor cell line and tumor-specific CTL are available in the laboratory, has been

Table III. Amino acid sequences of monoclonal TCR transcripts expressed in four parental and/or variant p53₂₆₄₋₂₇₂-specific CTL lines^a

Line	IVS	Specificity	V β Family	Sequences		
				V β	CDR3 region	J region
2	wt	wt + 7W	V β 13.6	RLELAAPSQTSVYFCA	SSQTPLG	DTQYFGPGTRLT/BJ2-3
4	wt	wt + 7W	V β 1	LELGDSALYFCA	SSEGGL	ETQYFGPGTRL/BJ2-5
53	7W	7W	V β 9	LGDSAVYFCA	SSAGTNT	YEQYFGPGTRLT/BJ2-7
68	7W	wt + 7W	V β 13.6	RLELAAPSQTSVYFCA	SSQTPLG	DTQYFGPGTRLT/BJ2-3

^a Lines 2 and 4 were stimulated with the parental wt p53₂₆₄₋₂₇₂ peptide (wt) and were reactive against the parental and 7W variant peptides. Lines 53 and 68 were stimulated with the 7W variant peptide. Line 53 was reactive against the 7W variant peptide only, while line 68 was reactive against the parental and 7W variant peptides. Single peaks in individual TCR variable chain families, suggesting clonality, were analyzed by direct sequencing of the PCR products.

Table IV. Summary of evidence of limited TCR V β usage for CTL recognition of the wt p53₂₆₄₋₂₇₂ epitope^a

PBMC Donors ^b	Induced with p53 ₂₆₄₋₂₇₂ Peptide	Bulk T Cell Population V β Usage	Derived T Cell Lines ^c	
			V β usage	p53 peptide specificity
Normal donor 2 (R)	wt	V β 1, 13.6	4 V β 1	wt/7W
Normal donor 4 ^d (R)	wt	V β 1	ND	
Normal donor 7 (NR)	7W	V β 9, 13.6	53 V β 9	7W only
			68 V β 13.6 ^e	wt/7W
OSCC patient 2 ^f (R)	wt	V β 13.6	2 V β 13.6 ^e	wt/7W

^a See Table III for the details on TCR V β usage.

^b Normal donors and patients identified in Tables I and II; R, responsive to IVS of PBMC to the wt p53₂₆₄₋₂₇₂ peptide; NR, nonresponsiveness.

^c T cell lines derived by limiting dilution.

^d Normal donor used as source of PBMC for induction of a bulk population of CTL specific for wt p53₂₆₄₋₂₇₂. Generation and characterization of this cell line was detailed in Ref. 7.

^e These T cell lines express identical V β , CDR3, and J region sequences (see Table III).

^f The OSCC patient 2 identified in Table I in Ref. 12.

of particular interest. The tumor cell line established from this patient and designated PCI-13 accumulates p53 molecules expressing a missense mutation at codon 286 and naturally presents the p53₂₆₄₋₂₇₂ epitope, albeit following pretreatment with IFN- γ (7, 12). The ability to generate anti-p53₂₆₄₋₂₇₂ CTL with the 7W variant from this patient's PBMC, which were nonresponsive to the parental peptide, provides a basis for the potential use of the 7W variant peptide in immunotherapy of this patient and, perhaps, other nonresponsive OSCC patients with tumors expressing similar characteristics. Concurrently, it needs to be determined whether the trends observed in OSCC patients regarding their responsiveness to wt p53 epitopes and the potential of their tumors to present these epitopes are also apparent in patients with other types of cancers.

References

- Houbiers, J. G. A., H. W. Nijman, J. W. Drijfhout, P. Kenemans, C. J. H. van der Velde, A. Brand, F. Momburg, W. M. Kast, and C. J. M. Melief. 1993. In vitro induction of human cytotoxic T lymphocyte responses against peptides of mutant and wild-type p53. *Eur. J. Immunol.* 23:2072.
- Theobald, M., J. Biggs, D. Dittmer, A. J. Levine, and L. A. Sherman. 1995. Targeting p53 as a general tumor antigen. *Proc. Natl. Acad. Sci. USA* 92: 11993.
- Mayordomo, J. I., D. J. Loftus, H. Sakamoto, C. M. DeCesare, P. M. Appasamy, M. T. Lotze, W. J. Storkus, E. A. Appella, and A. B. DeLeo. 1996. Therapy of murine tumors with p53 wild-type and mutant sequence peptide-based vaccines. *J. Exp. Med.* 183:1357.
- DeLeo, A. B. 1998. p53-based immunotherapy of cancer. *Crit. Rev. Immunol.* 18:29.
- Vierboom, M. P. M., D. I. Gabrilovich, R. Offringa, W. M. Kast, and C. J. M. Melief. 2000. p53: a target for T-cell mediated immunotherapy. In *Peptide-Based Cancer Vaccines*. W. M. Kast, ed. Landes Bioscience, Georgetown, p. 40.
- Ropke, M., J. Hald, P. Guldborg, J. Zeuthen, J. Norgaard, L. Fugger, A. Svejgaard, S. van der Burg, H. W. Nijman, C. J. M. Melief, and M. H. Claesson. 1996. Spontaneous human squamous cell carcinomas are killed by a human cytotoxic T lymphocyte clone recognizing a wild-type p53-derived peptide. *Proc. Natl. Acad. Sci. USA* 93:14704.
- Chikamatsu, K., K. Nakano, W. J. Storkus, E. Appella, M. T. Lotze, T. L. Whiteside, and A. B. DeLeo. 1999. Generation of anti-p53 cytotoxic T lymphocytes from human peripheral blood using autologous dendritic cells. *Clin. Cancer Res.* 5:1281.
- Gnjatic, S., Z. Cai, M. Viguier, S. Chouaib, J.-G. Guillet, and J. Chopin. 1998. Accumulation of the p53 protein allows recognition by human CTL of a wild-type p53 epitope presented by breast carcinomas and melanomas. *J. Immunol.* 160:328.
- Barfoed, A. M., T. R. Petersen, A. F. Kirkin, P. Thor Straten, M. H. Claesson, and J. Zeuthen. 2000. Cytotoxic T-lymphocyte clones, established by stimulation with the HLA-A2 binding p53₆₅₋₇₃ wild type peptide loaded on dendritic cells in vitro, specifically recognize and lyse HLA-A2 tumor cells overexpressing the p53 protein. *Scand. J. Immunol.* 51:128.
- McArdle, S. E. B., R. C. Rees, K. A. Mulcahy, J. Saba, C. A. McIntyre, and A. K. Murray. 2000. Induction of human cytotoxic T lymphocytes that preferentially recognize tumor cells bearing a conformational p53 mutant. *Cancer Immunol. Immunother.* 49:417.
- Eura, M., K. Chikamatsu, F. Katsura, A. Obata, Y. Sabao, M. Takiguchi, Y. Song, E. Appella, T. L. Whiteside, and A. B. DeLeo. 2000. A wild type-sequence p53 peptide presented by HLA-A24 induces cytotoxic T lymphocytes that recognize squamous cell carcinomas of the head and neck. *Clin. Cancer Res.* 6:979.
- Hoffmann, T. K., K. Nakano, E. Elder, G. Dworacki, S. D. Finkelstein, E. Appella, T. L. Whiteside, and A. B. DeLeo. 2000. Generation of T cells specific for the wild-type sequence p53₂₆₄₋₂₇₂ peptide in cancer patients: implications for immunoselection of epitope-loss variants. *J. Immunol.* 165:5938.
- Theobald, M., J. Biggs, J. Hernandez, J. Lustgarten, C. Labadie, and L. A. Sherman. 1997. Tolerance to p53 by A2.1-restricted cytotoxic T lymphocytes. *J. Exp. Med.* 185:833.
- Hernandez, J., P. L. Lee, M. M. Davis, and L. A. Sherman. 2000. The use of HLA-A2.1/p53 peptide tetramers to visualize the impact of self-tolerance on the TCR repertoire. *J. Immunol.* 164:596.
- Hernandez, J., A. Ko, and L. A. Sherman. 2001. CTLA-4 blockage enhances the CTL responses to the p53 self-tumor antigen. *J. Immunol.* 166:3908.
- Theobald, M., T. Ruppert, U. Kuckelkorn, J. Hernandez, A. Häussler, E. A. Ferreira, U. Liewer, J. Biggs, A. J. Levine, C. Huber, et al. 1998. The sequence alteration associated with a mutational hotspot in p53 protects cells from lysis by cytotoxic T lymphocytes specific for a flanking peptide epitope. *J. Exp. Med.* 188:1017.
- Boehnke, W. H., T. Takeshita, C. D. Pendleton, S. Sadegh-Nasseri, L. Racioppi, R. A. Houghton, J. A. Berzofsky, and R. N. Germain. 1993. The importance of dominant negative effects of amino acids side chain substitution in peptide-MHC molecule interactions and T cell recognition. *J. Immunol.* 150:331.
- Zaremba, S., E. Barzaga, M. Zhu, N. Soares, K. Y. Tsang, and J. Scholm. 1997. Identification of an enhancer agonist cytotoxic T lymphocytes peptide from human carcinoembryonic antigen. *Cancer Res.* 57:4570.
- Loftus, D. J., C. Castelli, T. M. Clay, P. Squarcina, F. M. Marincola, M. I. Nishimura, G. Parmiani, E. Appella, and L. Rivoltini. 1996. Identification of epitope mimics recognized by CTL reactive to the melanoma/melanocyte-derived peptide MART-1₂₇₋₃₅. *J. Exp. Med.* 184:647.
- Loftus, D., P. Squarcina, N. Mai-Britt, C. Geisler, C. Castelli, N. Odum, E. Appella, G. Parmiani, and L. Rivoltini. 1998. Peptides derived from self-proteins as partial agonists and antagonists of human CD8⁺ T-cell clones reactive to melanoma/melanocyte epitope MART-1₂₇₋₃₅ elicits anti-melanoma CD8⁺ T cells with enhanced functional characteristics: implication for more effective immunotherapy. *Cancer Res.* 58:2433.
- Rivoltini, L., P. Squarcina, D. Loftus, C. Castelli, P. Tarsini, A. Mazzocchi, F. Rini, V. Viggiano, F. Belli, and G. Parmiani. 1999. A superagonist variant of peptide MART1/Melan A₂₇₋₃₅ elicits anti-melanoma CD8⁺ T cells with enhanced functional characteristics: implication for more effective immunotherapy. *Cancer Res.* 59:301.
- Irvine, K. R., M. R. Parkhurst, E. P. Shulman, J. P. Tupesis, M. Custer, C. E. Touloukian, P. F. Robbins, A. G. Yafal, P. Greenhalgh, R. P. M. Suttmuller, et al. 1999. Recombinant virus vaccination against "self" antigens using anchor fixed immunogens. *Cancer Res.* 59:2536.
- Slansky, J. E., F. M. Rattis, L. F. Boyd, T. Fahmy, E. M. Jaffee, J. P. Schneck, D. H. Margulies, and D. M. Pardoll. 2000. Enhanced antigen-specific antitumor immunity with altered peptide ligands that stabilize the MHC-peptide-TCR complex. *Immunity* 13:529.
- Kalergis, A. M., N. Boucheron, M.-A. Doucey, E. Palmieri, E. C. Goyarts, Z. Vegh, E. F. Luescher, and S. G. Nathenson. 2001. Efficient T cell activation requires an optimal dwell-time of interaction between the TCR and the pMHC complex. *Nat. Immun.* 2:229.
- Dyall, R., W. B. Bowne, L. W. Weber, J. LeMaout, P. Szabo, Y. Moroi, G. Piskun, J. J. Lewis, A. N. Houghton, and J. Nikolic-Zugic. 1998. Heteroclitic immunization induces tumor immunity. *J. Exp. Med.* 188:1553.
- Petersen, T. R., S. Buus, S. Brunak, M. H. Nissen, L. A. Sherman, and M. H. Claesson. 2001. Identification and design of p53-derived HLA-A2-binding peptides with increased immunogenicity. *Scand. J. Immunol.* 53:357.

27. Heo, D. S., C. Snyderman, S. M. Gollin, S. Pan, P. Walker, R. Deka, E. L. Barnes, R. B. Herberman, and T. L. Whiteside. 1989. Biology, cytogenetics, and sensitivity to immunological effector cells of new head and neck squamous cell carcinoma lines. *Cancer Res.* 49:5167.
28. Sallusto, F., and A. Lanzavecchia. 1994. Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin-4 and down-regulated by tumor necrosis factor α . *J. Exp. Med.* 179:1109.
29. Hoffmann, T. K., N. Meidenbauer, G. Dworacki, H. Kanaya, and T. L. Whiteside. 2000. Generation of tumor-specific T lymphocytes by cross-priming with human dendritic cells ingesting apoptotic tumor cells. *Cancer Res.* 60:3542.
30. Hoffmann, T. K., V. Donnenberg, U. Friebe, M. Meyer, C. R. Rinaldo, A. B. DeLeo, T. L. Whiteside, and A. D. Donnenberg. 2000. Competition of peptide-MHC class I tetrameric complexes with anti-CD3 provides evidence for specificity of peptide binding to the TCR complex. *Cytometry* 41:3.21.
31. Puisieux, I., J. Even, C. Pannetier, F. Jotereau, M. Favrot, and P. Kourilsky. 1994. Oligoclonality of tumor infiltrating lymphocytes from human melanomas. *J. Immunol.* 153:2807.
32. Jager, E., M. Maeurer, H. Hohn, J. Karbach, D. Jager, Z. Zikianakis, A. Bakhshandeh-Bath, J. Orth, C. Neukirch, A. Necker, et al. 2000. Clonal expansion of Melan A-specific cytotoxic T lymphocytes in a melanoma patient responding to continued immunization with melanoma-associated peptides. *Int. J. Cancer* 86:538.
33. Monsurro, V., M.-B. Nielsen, A. Perez-Diez, M. E. Dudley, E. Wang, S. A. Rosenberg, and F. M. Marincola. 2001. Kinetics of TCR use in response to repeated epitope-specific immunization. *J. Immunol.* 166:5817.
34. Nikitina, E. Y., J. I. Clark, J. van Beynen, S. Chada, A. K. Virmani, D. P. Carbone, and D. I. Gabrilovich. 2001. Dendritic cells transduced with full-length wild-type p53 generate antitumor cytotoxic T lymphocytes from peripheral blood of cancer patients. *Clin. Cancer Res.* 7:127.